

# Enhanced Coastal Mapping Using Lidar Waveform Features

Christopher Parrish, Jeffrey Rogers, Larry Ward,  
and Jennifer Dijkstra

15th Annual JALBTCX

Airborne Coastal Mapping & Charting Workshop


10-12 June 2014



National Oceanic and Atmospheric Administration




# Recall from last year's JALBTCX presentation: *Simple, shape-based waveform features*



## Questions

- Are there simple, shape-based features that characterize the waveforms?
- Can they be computed in real-time?
- Can they be gridded and ingested into GIS?
- **Are they useful?**


# Animation of waveforms in transect across a marsh



# Waveform Features & Computation Times

Symbol	Metric Name	Computation Time ( $\mu\text{s}$ )
$w$	Width	14.8
$A$	Amplitude	0.7
$w/A$	Pulse aspect ratio	17.6
$AUC$	Area under curve	0.8
$AUC_r$	Area under curve: R/L ratio	17.8
$\beta_t$	Slope trailing edge	37.9
$\beta_r$	Slope ratio	38.4
$\sigma_w$	Standard deviation	0.9
$\mu_w$	Mean	0.7
$n_{50}$	Median	9.3
$\hat{n}$	Mode	0.6
$\gamma_1$	Skewness	8.2
$\beta_2$	Kurtosis	8.2
$I_1$	Pearson's 1st skewness coefficient	7.2
$I_2$	Pearson's 2nd skewness coefficient	12.3
$R_G^2$	Goodness-of-fit of Gaussian	850.3

# Gridded waveform features in GIS





# Results of regressions of $\Delta Z$ on waveform metrics ( $R^2$ )

<b>Metric</b>	<b>Little Pamet</b>	<b>Great Island</b>	<b>Moors</b>	<b>Mean</b>
Width	0.55	0.72	0.53	0.60
Standard deviation	0.55	0.73	0.42	0.57
Mean	0.27	0.27	0.04	0.19
Median	0.27	0.28	0.04	0.20
Mode	0.24	0.29	0.02	0.18
Goodness-of-fit of Gaussian	0.15	0.54	0.01	0.23


<b>Regression</b>	<b>Little Pamet</b>	<b>Great Island</b>	<b>Moors</b>	<b>Mean</b>
$\Delta Z$ on width and median	0.55	0.75	0.54	0.61
$\Delta Z$ on width and mean	0.55	0.75	0.53	0.61
$\Delta Z$ on width and mode	0.55	0.74	0.54	0.61
$\Delta Z$ on width and goodness-of-fit of Gaussian	0.56	0.80	0.56	0.64
$\Delta Z$ on PC1 and PC2	0.55	0.73	0.53	0.60

# Generating Relative Uncertainty Surfaces from Waveform Features



# Pamet marsh

## Relative uncertainty surface




Circles = field sample sites  
Blue = TF *Spartina alterniflora*




National Oceanic and Atmospheric Administration


# Use case #2 of lidar waveform features: Predicting salt marsh vegetation biophysical parameters



- = *S. alterniflora* samples
- = all other species







# Predicting salt marsh vegetation biophysical parameters

Results of correlations ( $r$ ) of biophysical parameters on waveform metrics for all vegetation species and the subset of *S. alterniflora*. Gray shaded cells have a  $p$  value  $<0.05$  ( $df = 24$ ).

Parameters	Width		Sample Skewness		Amplitude		Waveform Standard deviation		Pearson's 1st Skewness	
	All	<i>S. alterniflora</i>	All	<i>S. alterniflora</i>	All	<i>S. alterniflora</i>	All	<i>S. alterniflora</i>	All	<i>S. alterniflora</i>
Photographic Vegetation Height	<b>0.82</b>	<b>0.75</b>	<b>0.54</b>	0.17	<b>0.57</b>	0.17	<b>0.73</b>	<b>0.78</b>	0.36	0.37
Planimetric Obscuration	<b>0.47</b>	0.14	<b>0.56</b>	0.33	<b>0.71</b>	<b>0.62</b>	0.10	0.30	0.10	0.20
Quadrat Stem Density	<b>0.58</b>	<b>0.66</b>	<b>0.63</b>	0.35	<b>0.73</b>	0.00	0.35	<b>0.48</b>	<b>0.39</b>	0.00
Quadrat Biomass Density	<b>0.41</b>	0.20	0.30	0.22	<b>0.53</b>	0.14	0.17	0.22	0.00	0.14
Proportion Vegetation Area (25cm)	<b>0.73</b>	<b>0.57</b>	<b>0.45</b>	0.00	<b>0.39</b>	<b>0.61</b>	<b>0.62</b>	<b>0.46</b>	0.37	0.24
Proportion Vegetation Area (10cm)	<b>0.49</b>	0.14	0.28	0.14	0.26	0.28	0.33	0.00	0.22	0.00



National Oceanic and Atmospheric Administration

# Predicting salt marsh vegetation biophysical parameters

Results of multiple linear regressions ( $R^2$ ) of biophysical parameter with waveform metrics. Bold with underline represent improved results.

	Waveform Width and Sample Skewness	Waveform Width and Amplitude	Waveform Width and Waveform STDV	Sample Skewness and Amplitude	Sample Skewness and Waveform STDV	Amplitude and Waveform STDV
Vegetation Height	0.68	<b><u>0.72</u></b>	0.68	0.38	0.57	<b><u>0.74</u></b>
Planimetric Obscuration	0.32	0.53	0.49	0.53	0.36	0.51
Quadrat Stem Density	0.47	<b><u>0.6</u></b>	0.38	0.58	0.42	<b><u>0.6</u></b>
Quadrat Biomass Density	0.18	0.32	0.27	0.32	0.05	0.29
PVA (25 cm)	0.54	0.54	0.54	0.23	0.42	0.47




# Can we extend this to topo-bathy lidar and benthic habitat mapping?

DeHavilland Twin Otter  
(DHC-6)



Sept 2013 NOAA/NGS data of Barnegat Inlet



Left: Riegl LMS Q-680,  
Right: Riegl VQ-820-G



# Riegl waveform features

- Waveform features included as standard output from “V-line” systems and provided via LAS ExtraBytes

## 1. “Reflectance”

- Ratio of signal amplitude to amplitude of signal from a white reference target at same range, given in dB

$$A_{dB} = 10 \cdot \log_{10} \left( \frac{P_{echo}}{P_{DL}} \right)$$

$$\rho = A_{dB} - A_{dB,ref}(R)$$

## 2. Pulse Shape Deviation

- Measure of the discrepancy between the digitized waveform  $y[n]$  and a stored, system-specific reference pulse,  $p[n]$

$$\delta = \sum_{n=0}^{N-1} |y[n] - p[n]|$$




# Pre-Processing Steps

- No rigorous radiometric calibration (e.g., inversion of radiative transfer model) to solve for true bottom reflectance
- Instead, we apply some simple radiometric balancing to remove salient artifacts in mosaics of  $\rho_{rel}$  (or other gridded waveform features)
- Procedure
  - For each flightline and each waveform feature, compute the mean,  $\mu_i$ , and standard deviation,  $\sigma_i$
  - Pick one flightline that has good contrast and average “brightness” to be the reference
  - Normalize histograms of other flightlines, as follows

$$r' = \frac{\sigma_{ref}}{\sigma_i} (r - \mu_i) + \mu_{ref}$$



# Example: Preprocessed “reflectance” layer




National Oceanic and Atmospheric Administration

# Something else you can do...

- Remove any remaining artifacts (e.g., seamlines between swaths) from waveform feature mosaics in the frequency domain using ERDAS Imagine


Input



- $f(x, y) \xrightarrow{\mathcal{F}} F(u, v)$
- Notch filter to remove frequency components corresponding to seamlines
- $F_c(u, v) \xrightarrow{\mathcal{F}^{-1}} f_c(x, y)$




Output




# Data Layers




Aerial RGB  
Image



Bathymetry




Pulse Shape  
Deviation



Reflectance



# Benthic Habitats



# Seagrass

## KEY INDICATORS

Water quality

Ecosystem health

Essential fish and shellfish habitat





Eelgrass  
*Zostera marina*



Widgeongrass  
*Ruppia maritima*


# Barnegat Bay Field Campaign: October 2013



National Oceanic and Atmospheric Administration

# Sand and Macroalgae


Camera Photo




Aerial RGB Image




Pulse Shape Deviation



Reflectance Image




Bathymetry



# Sand and Eelgrass


Camera Photo




Aerial RGB Image




Pulse Shape Deviation



Reflectance Image



Bathymetry



# Eelgrass


Camera Photo




Aerial RGB Image




Pulse Shape Deviation



Reflectance Image



Bathymetry



# Sand


Camera Photo




Aerial RGB Image




Pulse Shape Deviation




Reflectance Image



Bathymetry



# Barnegat Bay Preliminary Habitat Map



Nati



# Next Steps & Future Direction

- Object-based classification of Barnegat Bay benthic habitats
  - eCog
  - Rule set based on texture of waveform features, depth, dist from shoreline
- EAARL-B / ALPS implementation
  - Great data set, acquired very shortly before and after Sandy
  - Pre- and post-Sandy => habitat change analysis
- (Jeff's dissertation work) Marsh elevation correction factors, computed as a function of waveform features, distance from shoreline, elevation relative to MHW
  - 2 more papers to be submitted to JCR SI



# References

Parrish, C.E., J.N. Rogers, and B.R. Calder, 2014. Assessment of Waveform Shape Features for Lidar Uncertainty Modeling in a Coastal Salt Marsh Environment. *Geoscience and Remote Sensing Letters*, Vol. 11, No. 2, pp. 569-573.

Pfennigbauer, M. and A. Ullrich, 2010. Improving quality of laser scanning data acquisition through calibrated amplitude and pulse deviation measurement. *Proc. SPIE Defense, Security, and Sensing*, pp. 76841F-76841F.

Rogers, J.N., C.E. Parrish, L. Ward, and D. Burdick, 2014. Evaluation of Vertical Obscuration and Full Waveform Lidar to Predict Salt Marsh Vegetation Biophysical Parameters. *Remote Sensing of Environment* (in revision).

+ 2 more papers to be submitted to JCR SI

